

Binary spinning black hole Hamiltonian in canonical center-of-mass and rest-frame coordinates through higher post-Newtonian order

Tilman J. Rothe

*Mathematisches Institut, Friedrich-Schiller-Universität Jena,
Ernst-Abbe-Platz 2, 07743 Jena, Germany*

Gerhard Schäfer

*Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena,
Max-Wien-Platz 1, 07743 Jena, Germany*

(Dated: March 1, 2010)

Abstract

The recently constructed Hamiltonians for spinless binary black holes through third post-Newtonian order and for spinning ones through formal second post-Newtonian order, where the spins are counted of zero post-Newtonian order, are transformed into fully canonical center-of-mass and rest-frame variables. The mixture terms in the Hamiltonians between center-of-mass and rest-frame variables are in accordance with the relation between the total linear momentum and the center-of-mass velocity as demanded by global Lorentz invariance. The various generating functions for the center-of-mass and rest-frame canonical variables are explicitly given in terms of the single-particle canonical variables. The no-interaction theorem does not apply because the world-line condition of Lorentz covariant position variables is not imposed.

PACS number(s): 11.30.Cp, 04.25.Nx, 45.20.Jj, 45.50.Pk

I. INTRODUCTION

The conservative Hamiltonians for non-spinning and spinning binary black holes are known to higher post-Newtonian (PN) orders in global inertial reference frames [1–9]. However, applications are typically made in the rest frame where the total linear momentum of the binary system vanishes [10–12]. In that case six degrees of freedom in phase space get suppressed and the remaining Hamiltonians simplify a lot. By boosting the system back, the original Hamiltonians are recovered but within a set of canonical variables different from the former single-particle ones. If a Hamiltonian were given in Lorentz covariant coordinates, boosting would be straightforward. However, the canonical coordinates the Hamiltonians are presented in are quite different from Lorentz covariant ones, so boosting of those coordinates is quite an involved procedure. In this paper we will solve the problem of boosted representation of Hamiltonians by constructing fully canonical center-of-mass and rest-frame coordinates. This work is based on the diploma thesis of one of the authors [13].

A historical view to the relativistic mechanics in Hamiltonian form, see, e.g. [14], clearly shows that the problem of constructing center-of-mass and rest-frame coordinates for interacting relativistic systems has a long tradition beginning with the pioneering works by Thomas, Bakamjian, and Foldy around the 1950s of the last century [15–17]. To our best knowledge, the known general relativistic gravitational Hamiltonians for binary point-like particles with and without spin have never been given in canonical center-of-mass and rest-frame coordinates, not even at the 1PN level where the world-line condition still applies, i.e. where fully canonical coordinates and Lorentz covariance of particle position vectors are still compatible, see, e.g. [18, 19]. Evidently, the center-of-mass and relative position coordinates we are searching for are of Newton-Wigner type because of component-wise vanishing Poisson brackets [20].

The importance of the present work can be seen in two directions (i), it is by far not trivial to explicitly construct the involved canonical coordinate transformations in phase space and (ii), for future applications, the ground will be led for a Hamiltonian treatment of e.g. the recoil in binary systems through gravitational radiation emission or of the orbital motion or scattering of binaries in many-body systems, see, e.g., [21] for the post-Newtonian motion of the Moon or [22] for a relativistic generalization of the Jacobi momenta of the non-relativistic three-body problem.

II. IMPLICATIONS FROM THE POINCARÉ ALGEBRA

In space-asymptotically flat spacetimes, the ten conserved quantities total energy H , linear momentum $P^i \equiv P_i$, angular momentum $J^i \equiv J_i$, and Lorentz boost vector K^i have to fulfill the Poincaré algebra

$$\{P^i, H\} = \{J^i, H\} = 0, \quad (1a)$$

$$\{G^i, H\} = P^i, \quad \{P^i, P^j\} = 0, \quad (1b)$$

$$\{J^i, P^j\} = \epsilon^{ijk} P^k, \quad \{J^i, J^j\} = \epsilon^{ijk} J^k, \quad (1c)$$

$$\{J^i, G^j\} = \epsilon^{ijk} G^k, \quad \{G^i, P^j\} = \frac{1}{c^2} H \delta^{ij}, \quad (1d)$$

$$\{G^i, G^j\} = -\frac{1}{c^2} \epsilon^{ijk} J^k, \quad (1e)$$

where $G^i = K^i + P^i t$ is the non-explicitly time-dependent center-of-energy vector G^i .

Defining

$$M = \frac{1}{c^2} \sqrt{H^2 - c^2 P^2}, \quad \text{or,} \quad H = \sqrt{c^4 M^2 + c^2 P^2}, \quad (2)$$

a canonical center-of-mass coordinate X^i can be introduced in the following manner [23–26]

$$\mathbf{X} = c^2 \frac{\mathbf{G}}{H} + \frac{1}{M(H + c^2 M)} \left(\mathbf{J} - c^2 \left(\frac{\mathbf{G}}{H} \times \mathbf{P} \right) \right) \times \mathbf{P}, \quad (3a)$$

where $\mathbf{G} = (G^i)$ etc. Reversely, the center-of-energy vector can be expressed in the form

$$\mathbf{G} = \frac{1}{c^2} H \mathbf{X} - \frac{1}{M c^2 + H} (\mathbf{J} - \mathbf{X} \times \mathbf{P}) \times \mathbf{P}. \quad (3b)$$

It is crucial to point out that we make no further assumptions on $\mathbf{X}$ or $\mathbf{G}$.

The fundamental Poisson brackets read

$$\{X^i, X^j\} = 0, \quad (4a)$$

$$\{X^i, P^j\} = \delta^{ij}, \quad (4b)$$

$$\{P^i, P^j\} = 0. \quad (4c)$$

Further, for the center-of-mass velocity $\mathbf{V}$,

$$\mathbf{V} \equiv \dot{\mathbf{X}} = \{\mathbf{X}, H\} = \frac{c^2 \mathbf{P}}{H}, \quad \text{or,} \quad \mathbf{P} = \frac{H}{c^2} \mathbf{V} \quad (5)$$

are valid. Hereof the relations

$$\mathbf{P} = \frac{M\mathbf{V}}{\sqrt{1 - \frac{V^2}{c^2}}} \quad \text{and} \quad H = \frac{Mc^2}{\sqrt{1 - \frac{V^2}{c^2}}} \quad (6)$$

follow as the single-particle case mandates. Notice that it also holds

$$\mathbf{V} = \{\mathbf{G}, H\} \frac{c^2}{H}, \quad (7)$$

i.e. the speed of the center-of-energy vector coincides with the speed of the canonical center-of-mass position vector.

The Poisson brackets resulting from the Poincaré algebra,

$$\{H, M\} = 0, \quad (8a)$$

$$\{P^i, M\} = 0, \quad (8b)$$

$$\{X^i, M\} = 0, \quad (8c)$$

are crucial in the following. They fully generally show that the rest-mass energy M is a constant of motion not depending on the center-of-mass phase-space coordinates $\mathbf{X}$ and $\mathbf{P}$.

Restricting ourselves at the beginning to spinless binary point-mass systems, the explicit expressions for the total linear and orbital angular momentum read

$$\mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2, \quad \mathbf{J} = \mathbf{x}_1 \times \mathbf{p}_1 + \mathbf{x}_2 \times \mathbf{p}_2, \quad (9)$$

where $\mathbf{x}_1, \mathbf{p}_1$ and $\mathbf{x}_2, \mathbf{p}_2$ denote the canonical position and momentum variables of the particles with labels 1, 2.

In the following we will also need a slight generalization of an infinitesimal canonical transformation. Let g be the generator, that is a smooth function on phase space. Define by

$$T_g f \equiv \exp(\{\cdot, g\}) f \quad (10a)$$

$$= f + \{f, g\} + \frac{1}{2} \{\{f, g\}, g\} + \frac{1}{6} \{\{\{f, g\}, g\}, g\} + \dots \quad (10b)$$

an operator on phase-space functions f . It is easy to check that T_g (i), preserves Poisson brackets and (ii), does not modify Kronecker deltas, and therefore is a (finite) canonical transformation. For our purposes, the generator will always be at least of order c^{-2} , and we are only interested in a weakly relativistic portion. Cutting off after the appropriate PN level, only finitely many brackets remain in the above formula, and T_g thereby turns into a generalized infinitesimal canonical transformation.

III. CENTER-OF-MASS AND REST-FRAME CANONICAL COORDINATES

The well-known canonical coordinates adapted to the Newtonian problem obey the usual canonical Poisson bracket relations exactly without reference to any PN cutoff:

$$\mathbf{X}^N = \frac{m_1}{m_1 + m_2} \mathbf{x}_1 + \frac{m_2}{m_1 + m_2} \mathbf{x}_2, \quad (11a)$$

$$\mathbf{P}^N = \mathbf{p}_1 + \mathbf{p}_2, \quad (11b)$$

$$\mathbf{x}^N = \mathbf{x}_1 - \mathbf{x}_2, \quad (11c)$$

$$\mathbf{p}^N = \frac{m_2}{m_1 + m_2} \mathbf{p}_1 - \frac{m_1}{m_1 + m_2} \mathbf{p}_2, \quad (11d)$$

with

$$\{X^{Ni}, P^{Nj}\} = \delta^{ij}, \quad \{x^{Ni}, p^{Nj}\} = \delta^{ij}, \quad \text{zero otherwise.} \quad (12)$$

A contact transformation will take $\mathbf{X}^N$ to the correct Newton-Wigner center-of-mass coordinate $\mathbf{X}$. In the spirit of Ref. [16], we impose

$$T_g \mathbf{X}^N = \mathbf{X}, \quad (13a)$$

$$\{\mathbf{P}, g\} = 0, \quad (13b)$$

$$\{\mathbf{J}, g\} = 0, \quad (13c)$$

$$g = O(c^{-2}) \quad (13d)$$

on the generating function g and find the new coordinates as

$$\mathbf{X} = T_g \mathbf{X}^N, \quad (14a)$$

$$\mathbf{P} = T_g \mathbf{P}^N, \quad (14b)$$

$$\mathbf{x} = T_g \mathbf{x}^N, \quad (14c)$$

$$\mathbf{p} = T_g \mathbf{p}^N. \quad (14d)$$

These conditions have a number of desirable consequences. The new set of fundamental canonical coordinates includes the center-of-mass linear momentum and position vectors $\mathbf{P}$ and $\mathbf{X}$, respectively. Poisson bracket relations similar to the particle variables are satisfied, no constraints are necessary. The orbital angular momentum is realized simply as $\mathbf{J} = \mathbf{X} \times \mathbf{P} + \mathbf{x} \times \mathbf{p}$. They also entail that the center-of-energy vector $\mathbf{G} = \mathbf{G}(\mathbf{X}, \mathbf{P}, \mathbf{J}, M)$ depends only on the motion of the system as a whole, while the invariant mass $M = M(\mathbf{x}, \mathbf{p})$ is a function of the internal dynamics only. The latter is important because M generates

the internal part of the equations of motion: for any observable f , we have $\{f, H\} = c^2 M H^{-1} \{f, c^2 M\} + \frac{1}{2} c^2 H^{-1} \{f, P^2\}$. Since T_g is a canonical transformation, the Poincaré algebra is satisfied in the new variables.

The generator of the canonical transformation we calculated by the method of undetermined coefficients [2]. All algebraic manipulations were performed with the aid of MATHEMATICA and XTENSOR [27]. Groebner basis methods were used to account for the non-uniqueness resulting from the following vector identity:

$$(\mathbf{v}_1 \cdot \mathbf{v}_2 \times \mathbf{v}_3) \mathbf{v}_4 = (\mathbf{v}_4 \cdot \mathbf{v}_2 \times \mathbf{v}_3) \mathbf{v}_1 + (\mathbf{v}_1 \cdot \mathbf{v}_4 \times \mathbf{v}_3) \mathbf{v}_2 + (\mathbf{v}_1 \cdot \mathbf{v}_2 \times \mathbf{v}_4) \mathbf{v}_3. \quad (15)$$

IV. RESULTS FOR TWO POINT MASSES WITHOUT SPIN

It is most convenient to introduce rescaled variables in the form

$$\hat{\mathbf{x}} = \frac{\mathbf{x}}{Gm}, \quad \hat{\mathbf{p}} = \frac{\mathbf{p}}{\mu}, \quad \hat{t} = \frac{t}{Gm}, \quad \hat{M} = \frac{M - m}{\mu}, \quad (16a)$$

$$\hat{r} = |\hat{\mathbf{x}}|, \quad \hat{\mathbf{n}} = \hat{\mathbf{x}}/\hat{r} = \mathbf{x}/r = \mathbf{n}, \quad (16b)$$

$$\mu = \frac{m_1 m_2}{m_1 + m_2}, \quad m = m_1 + m_2, \quad \nu = \frac{\mu}{m}. \quad (16c)$$

From the structure of our Hamiltonian in relation to the rest-mass energy it is clear that our reduced rest mass has to read [10, Eqs. 3.2-3.6], [11], [4],

$$\begin{aligned} \hat{M} = & \frac{1}{2} \hat{p}^2 - \frac{1}{\hat{r}} + c^{-2} \left\{ \frac{3\nu - 1}{8} (\hat{p}^2)^2 - \frac{\nu + 3}{2} \frac{\hat{p}^2}{\hat{r}} - \frac{\nu}{2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2}{\hat{r}} + \frac{1}{2} \frac{1}{\hat{r}^2} \right\} \\ & + c^{-4} \left\{ \frac{1}{16} (5\nu^2 - 5\nu + 1) (\hat{p}^2)^3 - \frac{3\nu^2 + 20\nu - 5}{8} \frac{(\hat{p}^2)^2}{\hat{r}} - \frac{\nu^2}{4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 \hat{p}^2}{\hat{r}} \right. \\ & \quad \left. - \frac{3\nu^2}{8} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^4}{\hat{r}} + \left(4\nu + \frac{5}{2} \right) \frac{\hat{p}^2}{\hat{r}^2} + \frac{3\nu}{2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2}{\hat{r}^2} - \frac{3\nu + 1}{4} \frac{1}{\hat{r}^3} \right\} \\ & + c^{-6} \left\{ \frac{5}{128} (7\nu^3 - 14\nu^2 + 7\nu - 1) (\hat{p}^2)^4 + \frac{1}{16} (-5\nu^3 - 53\nu^2 + 42\nu - 7) \frac{(\hat{p}^2)^3}{\hat{r}} \right. \\ & \quad + \frac{(2 - 3\nu)\nu^2}{16} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 (\hat{p}^2)^2}{\hat{r}} - \frac{3(\nu - 1)\nu^2}{16} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^4 \hat{p}^2}{\hat{r}} - \frac{5\nu^3}{16} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^6}{\hat{r}} \\ & \quad + \frac{109\nu^2 + 136\nu - 27}{16} \frac{(\hat{p}^2)^2}{\hat{r}^2} + \frac{\nu(30\nu + 17)}{16} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 \hat{p}^2}{\hat{r}^2} \\ & \quad + \frac{\nu(43\nu + 5)}{12} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^4}{\hat{r}^2} - \frac{552\nu^2 + (1340 - 3\pi^2)\nu + 600}{192} \frac{\hat{p}^2}{\hat{r}^3} \\ & \quad \left. - \frac{\nu(112\nu + 3\pi^2 + 340)}{64} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2}{\hat{r}^3} + \frac{(872 - 63\pi^2)\nu + 12}{96} \frac{1}{\hat{r}^4} \right\} + O(c^{-8}), \end{aligned} \quad (17)$$

and the non-center-of-mass Hamiltonian has to be given by

$$H = \sqrt{c^4 \left(\mu \hat{M} + m \right)^2 + c^2 P^2}. \quad (18)$$

The generating function g_{point} as detailed by Eqs. (10) then turns out to read (note we are working with four different mass expressions m_1, m_2, m, μ for two different point masses to cut down on formula length), where $r_{12} = |\mathbf{x}_1 - \mathbf{x}_2|$, $\mathbf{n}_{12} = (\mathbf{x}_1 - \mathbf{x}_2)/r_{12}$,

$$g_{\text{point}} = \frac{1}{2} \left[g_{1\text{pN}} + g_{2\text{pN}} + g_{3\text{pN}}^{0\text{G}} + g_{3\text{pN}}^{1\text{G}} + g_{3\text{pN}}^{2\text{G}} + g_{3\text{pN}}^{3\text{G}} + (1 \leftrightarrow 2) \right], \quad (19)$$

$$g_{1\text{pN}} = c^{-2} \left\{ \left[\frac{m_2^2}{m_1 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_1^2 r_{12} + \frac{m_2^2}{m_1 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_2) p_1^2 r_{12} - \frac{2m_2 - m}{m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2) r_{12} \right] - G \frac{\mu(2m_2 - m)}{m} (\mathbf{n}_{12} \cdot \mathbf{p}_1) \right\}, \quad (20a)$$

$$\begin{aligned} g_{2\text{pN}} = c^{-4} \left\{ \left[- \frac{m_2^2(m_2 + 2m_1)}{4m_1^3 m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (p_1^2)^2 r_{12} - \frac{m_2^2(m_2 + 2m_1)}{4m_1^3 m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (p_1^2)^2 r_{12} \right. \right. \\ + \frac{2m_2 + \mu - m}{2m_1^2 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2) p_1^2 r_{12} + \frac{2m_2 + \mu - m}{2m_1^2 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{p}_1 \cdot \mathbf{p}_2) p_1^2 r_{12} \\ - \frac{2m_2 - m}{2\mu m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2)^2 r_{12} - \frac{2m_2 - m}{4\mu m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_2^2 p_1^2 r_{12} \left. \right] \\ - G \left[\frac{m_2^2(m_2 - m_1)}{4m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 + \frac{m_1(-5m_2 + 6\mu + m)}{4m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 \right. \\ + \frac{2\mu(3\mu - 8m) - 5m_2(\mu - 2m)}{4m_1 m^2} (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_1^2 \\ + \frac{6\mu^2 - 5m_2\mu - 16m\mu + 5m^2 + 5m_2 m}{4m_1 m^2} (\mathbf{n}_{12} \cdot \mathbf{p}_2) p_1^2 \\ + \frac{6m^2 - 17m_2 m - 3\mu m + 6m_2\mu}{2m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2) \left. \right] \\ + G^2 \frac{\mu(2\mu - m)(m - 2m_2)}{2m} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)}{r_{12}} \left. \right\}, \quad (20b) \end{aligned}$$

$$\begin{aligned}
g_{3\text{pN}}^{0\text{G}} = c^{-6} & \left\{ \frac{m_2(-\mu^2 + 6m^2) + 2\mu(\mu^2 + 5m\mu - 3m^2)}{48m_1^5m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (p_1^2)^3 r_{12} \right. \\
& + \frac{m_2(-\mu^2 + 6m^2) + 2\mu(\mu^2 + 5m\mu - 3m^2)}{48m_1^5m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (p_1^2)^3 r_{12} \\
& - \frac{m(3\mu^2 + 16m\mu - 8m^2) + m_2(-6\mu^2 - 20m\mu + 17m^2)}{24m_1^4m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2) (p_1^2)^2 r_{12} \\
& - \frac{m(3\mu^2 + 16m\mu - 8m^2) + m_2(-6\mu^2 - 20m\mu + 17m^2)}{24m_1^4m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{p}_1 \cdot \mathbf{p}_2) (p_1^2)^2 r_{12} \\
& + \frac{m_2(-6\mu^2 - 14m\mu + 13m^2) + m(3\mu^2 + 7m\mu - 6m^2)}{12m_2m_1^3m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2)^2 p_1^2 r_{12} \\
& - \frac{m_2(6\mu^2 + 14m\mu - 13m^2) + m(-3\mu^2 - 7m\mu + 6m^2)}{12m_2m_1^3m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{p}_1 \cdot \mathbf{p}_2)^2 p_1^2 r_{12} \\
& + \frac{(2m_2 - m)(\mu^2 + 2m\mu - 2m^2)}{6\mu^2m^7} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2)^3 r_{12} \\
& - \frac{2m_2(3\mu^2 + 7m\mu - 6m^2) + m(-3\mu^2 - 16m\mu + 7m^2)}{48m_2m_1^3m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_2^2 (p_1^2)^2 r_{12} \\
& - \frac{2m_2(3\mu^2 + 7m\mu - 6m^2) + m(-3\mu^2 - 16m\mu + 7m^2)}{48m_2m_1^3m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_2) p_2^2 (p_1^2)^2 r_{12} \\
& \left. + \frac{(2m_2 - m)(3\mu^2 + 6m\mu - 5m^2)}{12\mu^2m^7} (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_2^2 (\mathbf{p}_1 \cdot \mathbf{p}_2) p_1^2 r_{12} \right\}, \tag{20c}
\end{aligned}$$

$$\begin{aligned}
g_{3\text{pN}}^{1\text{G}} = & c^{-6}G \left\{ \frac{m_2^2(m_1 - m_2)}{16m^6} (\mathbf{n}_{12} \cdot \mathbf{p}_1)^5 + \frac{10 \mu^2 - m_2\mu - 16m\mu + m^2 + 5m_2m}{16m_1 m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)^4 \right. \\
& + \frac{10 \mu^2 + 3m_2\mu - 26m\mu + 8m^2 - 5m_2m}{8m_1 m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2 (\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 \\
& + \frac{m_2^2(10m_2 + 4\mu - 3m)}{48m_1 m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 p_1^2 \\
& - \frac{4\mu^2 - 13 m\mu + m^2 + 6m_2(\mu + m)}{16m_1 m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 p_1^2 \\
& + \frac{m(6\mu^2 + 9 m\mu - 4m^2) + m_2(-4\mu^2 + 15m\mu + 5 m^2)}{16\mu m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_2)^2 (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_1^2 \\
& - \frac{m_2(27\mu^2 - 32m\mu - 54 m^2) + 2\mu(\mu^2 - 9m\mu + 43m^2)}{48 m_1^3 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (p_1^2)^2 \\
& - \frac{m_2(27\mu^2 - 20m\mu - 42m^2) + 2(\mu^3 - 9m \mu^2 + 43m^2\mu - 6m^3)}{48m_1^3 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (p_1^2)^2 \\
& + \frac{-4m_2\mu + 2m\mu + 3m_2 m}{24m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 (\mathbf{p}_1 \cdot \mathbf{p}_2) \\
& - \frac{m(-2\mu^2 + 7m\mu - 6m^2) + m_2(4 \mu^2 - 15m\mu + 18m^2)}{8\mu m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2 (\mathbf{p}_1 \cdot \mathbf{p}_2) \\
& - \frac{m(-30\mu^2 + 84 m\mu - 47m^2) + m_2(4\mu^2 - 38m\mu + 101 m^2)}{24m_1^2 m^4} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2) p_1^2 \\
& - \frac{-4\mu^3 + 38m\mu^2 - 101m^2\mu - 6m^3 + m_2(-26\mu^2 + 22m\mu + 60m^2)}{24m_2 m_1^2 m^3} (\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{p}_1 \cdot \mathbf{p}_2) p_1^2 \\
& - \frac{m_2(-2\mu^2 + 14 m\mu - 55m^2) + m(\mu^2 - 13m\mu + 29 m^2)}{12\mu m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2)^2 \\
& - \frac{m_2(4\mu^2 - 11m\mu - 43m^2) + m(2\mu^2 + 13m\mu + 13m^2)}{48\mu m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1)^3 p_2^2 \\
& \left. - \frac{m_2(-4 \mu^2 + 60m\mu + 2m^2) + m(2\mu^2 - 42m \mu + 17m^2)}{48\mu m^5} (\mathbf{n}_{12} \cdot \mathbf{p}_1) p_2^2 p_1^2 \right\}, \tag{20d}
\end{aligned}$$

$$\begin{aligned}
g_{3\text{pN}}^{2\text{G}} = & c^{-6}G^2 \left\{ \frac{m_2(24\mu^2 - 20m\mu + 5m^2 + 2m_2(m - 6\mu))}{24m^3} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)^3}{r_{12}} \right. \\
& - \frac{m(12\mu^2 - 6m\mu - 5m^2) + m_2(-72\mu^2 + 48m\mu + 3m^2)}{24 m^3} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)^2}{r_{12}} \\
& + \frac{m_2(38\mu^2 + 19m\mu + 90m^2) - 2\mu(22 \mu^2 - 33m\mu + 68m^2)}{24m_1 m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) p_1^2}{r_{12}} \\
& + \frac{-44 \mu^3 + 66m\mu^2 - 166m^2\mu + 65m^3 + m_2(38 \mu^2 + 37m\mu + 25m^2)}{24m_1 m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2) p_1^2}{r_{12}} \\
& \left. - \frac{m(-44\mu^2 + 62m\mu - 79m^2) + 4m_2(22 \mu^2 - 40m\mu + 69m^2)}{24m^3} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2)}{r_{12}} \right\}, \tag{20e}
\end{aligned}$$

$$g_{3\text{pN}}^{3\text{G}} = c^{-6}G^3 \frac{\mu(m - 2m_2)(4\mu^2 - m\mu + m^2)}{4m} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1)}{r_{12}^2}. \tag{20f}$$

V. RESULTS FOR TWO PARTICLES WITH SPIN

In the case of two spinning particles, the Hamiltonian and the center-of-energy vector are only known up to the formal 2PN order. Here, we are counting formally, i.e. $\mathbf{S}_1$ has the same units as an orbital angular momentum, without any reference to its magnitude for maximally rotating black holes. Since the expression for H is quite lengthy, we will not repeat here the formulas for the Hamiltonians given in [6–9]. However, the total Hamiltonian may be abbreviated as follows, [28],

$$\begin{aligned}
H = & H_N + H_{1PN} + H_{2PN} + H_{3PN} \\
& + H_{SO}^{\text{LO}} + H_{S_1 S_2}^{\text{LO}} + H_{S_1^2}^{\text{LO}} + H_{S_2^2}^{\text{LO}} \\
& + H_{SO}^{\text{NLO}} + H_{S_1 S_2}^{\text{NLO}} + H_{S_1^2}^{\text{NLO}} + H_{S_2^2}^{\text{NLO}} \\
& + H_{p_1 S_2^3} + H_{p_2 S_1^3} + H_{p_1 S_1^3} + H_{p_2 S_2^3} \\
& + H_{p_1 S_1 S_2^2} + H_{p_2 S_2 S_1^2} + H_{p_1 S_2 S_1^2} + H_{p_2 S_1 S_2^2} \\
& + H_{S_1^2 S_2^2} + H_{S_1 S_2^3} + H_{S_2 S_1^3} ,
\end{aligned} \tag{21}$$

where LO and NLO respectively denote leading and next-to-leading order coupling and SO spin-orbit coupling. The other Hamiltonians with spin are leading order ones. The related expression for $\hat{M}$ is given by

$$\hat{M}_{2\text{pN spin}}^{\text{total}} = \hat{M}_{\text{point}} + \frac{1}{2} \left[\hat{M}_{1\text{pN}}^s + \hat{M}_{2\text{pN}}^s + (1 \leftrightarrow 2) \right] + O(c^{-6}) . \tag{22}$$

Here, “ $1 \leftrightarrow 2$ ” maps $\mathbf{p} \leftrightarrow -\mathbf{p}$, $\mathbf{x} \leftrightarrow -\mathbf{x}$, $\mathbf{S}_1 \leftrightarrow \mathbf{S}_2$, and $m_1 \leftrightarrow m_2$. In an abuse of notation, we will take $\hat{M}_{\text{point}}$ to signal the same dependency on the variables $\mathbf{x}$, $\mathbf{p}$ as in the spinless case. Since these have a different meaning here, $\hat{M}_{\text{point}}$ is not the same phase-space function as $\hat{M}$ in the previous section. Introducing

$$\hat{\mathbf{S}}_1 = \frac{\mathbf{S}_1}{Gm_1}, \quad \hat{\mathbf{S}}_2 = \frac{\mathbf{S}_2}{Gm_2}, \tag{23}$$

where

$$\mathbf{S}_1 = T_{g_{\text{spin}}}^{\text{total}} \mathbf{S}_1, \tag{24a}$$

$$\mathbf{S}_2 = T_{g_{\text{spin}}}^{\text{total}} \mathbf{S}_2, \tag{24b}$$

we get

$$\begin{aligned} \hat{M}_{1\text{pN}}^{\text{s}} = c^{-2} & \left\{ \frac{4m - m_2}{m^2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_1)}{r^2} + \frac{3}{m^2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^3} \right. \\ & \left. + \frac{3}{m^2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2)(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)}{r^3} - \frac{1}{m^2} \frac{\hat{\mathcal{S}}_1^2}{r^3} - \frac{1}{m^2} \frac{(\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)}{r^3} \right\}, \end{aligned} \quad (25\text{a})$$

$$\begin{aligned}
\hat{M}_{2\text{pN}}^s = c^{-4} & \left\{ \frac{2(m_2 - 2(\mu + 3m))}{m^2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_1)}{\hat{r}^3} + \frac{3m_2 m_1 (m_2 + 2m_1)}{2m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_1)}{\hat{r}^2} (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 \right. \\
& - \frac{3m_2 \mu - 19m\mu + 5m_2 m}{4m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_1) \hat{p}^2}{\hat{r}^2} - \frac{5m_2 + 9m}{m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^4} \\
& + \frac{15\mu}{2m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^3} - \frac{12}{m^2} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^4} \\
& + \frac{15\mu}{2m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^3} + \frac{3(-7m_2 + \mu + 6m)}{4m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2 \hat{p}^2}{\hat{r}^3} \\
& + \frac{9\mu + 6m}{4m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1) \hat{p}^2}{\hat{r}^3} - \frac{3(m_2 + \mu)}{2m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1) (\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^3} \\
& + \frac{3(2m_2 - 3\mu - 4m)}{2m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^3} + \frac{m_2^2}{2m^4} \frac{(\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^3} \\
& + \frac{\mu + 3m}{2m^3} \frac{(\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^3} + \frac{m_2 + 5m}{m^3} \frac{\hat{\mathcal{S}}_1^2}{\hat{r}^4} + \frac{3(m_2 - 3\mu)}{4m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 \hat{\mathcal{S}}_1^2}{\hat{r}^3} \\
& - \frac{3m_1 \hat{p}^2 \hat{\mathcal{S}}_1^2}{2m^3 \hat{r}^3} + \frac{6}{m^2} \frac{(\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)}{\hat{r}^4} - \frac{3(\mu - 4m)}{4m^3} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})^2 (\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)}{\hat{r}^3} \\
& - \frac{2\mu + 3m}{2m^3} \frac{\hat{p}^2 (\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)}{\hat{r}^3} - \frac{5(m_2 + 4m_1)}{2m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_1) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^4} \\
& + \frac{m_2 + 4m_1}{2m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_1) \hat{\mathcal{S}}_1^2}{\hat{r}^4} + \frac{15(m_2 - 3m)}{2m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^4} \\
& - \frac{3(m_2 - 3m)}{2m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}} \times \hat{\mathbf{S}}_2) \hat{\mathcal{S}}_1^2}{\hat{r}^4} - \frac{15m_2}{m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1 \times \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^4} \\
& + \frac{3m_2}{m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1 \times \hat{\mathbf{S}}_2) (\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^4} + \frac{6m_2 - 9m}{m^4} \frac{(\hat{\mathbf{p}} \cdot \hat{\mathbf{S}}_1 \times \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)}{\hat{r}^4} \\
& - \frac{35}{m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^3}{\hat{r}^5} - \frac{105}{4m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2)^2 (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2}{\hat{r}^5} \\
& + \frac{15}{m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1) \hat{\mathcal{S}}_1^2}{\hat{r}^5} + \frac{15}{2m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2)^2 \hat{\mathcal{S}}_1^2}{\hat{r}^5} \\
& + \frac{15}{m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1)^2 (\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)}{\hat{r}^5} + \frac{15}{m^4} \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_2) (\hat{\mathbf{n}} \cdot \hat{\mathbf{S}}_1) (\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)}{\hat{r}^5} \\
& \left. - \frac{3}{m^4} \frac{(\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2) \hat{\mathcal{S}}_1^2}{\hat{r}^5} + \frac{3}{4m^4} \frac{(\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2)^2}{\hat{r}^5} - \frac{3}{4m^4} \frac{\hat{\mathcal{S}}_2^2 \hat{\mathcal{S}}_1^2}{\hat{r}^5} \right\}. \tag{25b}
\end{aligned}$$

Again, we have determined a generating function such that the binary's invariant mass takes the same form in the transformed variables as in the center-of-mass frame. However, the Poincaré group generators H , $\mathbf{G}$, $\mathbf{J}$ and therefore also $\mathbf{X}$ differ from the point-particle case, particularly $\mathbf{J} = \mathbf{x}_1 \times \mathbf{p}_1 + \mathbf{x}_2 \times \mathbf{p}_2 + \mathbf{S}_1 + \mathbf{S}_2$, where additionally $\{S_a^i, S_b^j\} = \epsilon^{ijk} S^k \delta_{ab}$ holds.

The generating function $g_{\text{spin}}^{\text{total}}$ reads,

$$g_{\text{spin}}^{\text{total}} = g_{\text{point}} + \frac{1}{2} [g_{\text{spin}} + (1 \leftrightarrow 2)], \quad (26)$$

$$\begin{aligned} g_{\text{spin}} = & -c^{-2} \frac{1}{m_1 m} (\mathbf{p}_1 \cdot \mathbf{p}_2 \times \mathbf{S}_1) + c^{-4} \left\{ \left[\frac{\mu + m}{4m_1^3 m^2} (\mathbf{p}_1 \cdot \mathbf{p}_2 \times \mathbf{S}_1) p_1^2 \right. \right. \\ & + \frac{2m_2 - 3m}{4m_1^2 m^3} (\mathbf{p}_1 \cdot \mathbf{p}_2 \times \mathbf{S}_1) (\mathbf{p}_1 \cdot \mathbf{p}_2) - \frac{m_2 + m}{4\mu m^4} (\mathbf{p}_1 \cdot \mathbf{p}_2 \times \mathbf{S}_2) p_1^2 \Big] \\ & + G \left[\frac{3(m_2 - 2\mu)}{2m_1 m} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{S}_1)^2}{r_{12}^2} + \frac{6m_2 - 3m}{m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{S}_2) (\mathbf{n}_{12} \cdot \mathbf{S}_1)}{r_{12}^2} \right. \\ & + \frac{3(m_2 + 2\mu - m)}{2m_2 m} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{n}_{12} \cdot \mathbf{S}_2)^2}{r_{12}^2} - \frac{4m_2}{m_1 m} \frac{(\mathbf{n}_{12} \cdot \mathbf{S}_1) (\mathbf{p}_1 \cdot \mathbf{S}_1)}{r_{12}^2} \\ & + \frac{1}{m} \frac{(\mathbf{n}_{12} \cdot \mathbf{S}_2) (\mathbf{p}_1 \cdot \mathbf{S}_1)}{r_{12}^2} - \frac{1}{m} \frac{(\mathbf{n}_{12} \cdot \mathbf{S}_1) (\mathbf{p}_1 \cdot \mathbf{S}_2)}{r_{12}^2} \\ & + \frac{4m_1}{m_2 m} \frac{(\mathbf{n}_{12} \cdot \mathbf{S}_2) (\mathbf{p}_1 \cdot \mathbf{S}_2)}{r_{12}^2} + \frac{3m_2 + 2\mu}{2m_1 m} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) S_1^2}{r_{12}^2} \\ & + \frac{m - 2m_2}{m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) (\mathbf{S}_1 \cdot \mathbf{S}_2)}{r_{12}^2} - \frac{m_1(2m_2 + 3m)}{2m_2 m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1) S_2^2}{r_{12}^2} \\ & + \frac{2m_2(\mu + 5m) - 17\mu m}{4m_1 m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1 \times \mathbf{S}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_1)}{r_{12}} \\ & + \frac{4m^2 + 6m_2 m - 17\mu m + 2m_2 \mu}{4m_1 m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1 \times \mathbf{S}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_2)}{r_{12}} \\ & + \frac{19m_2 - 2\mu - 7m}{4m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_1 \times \mathbf{S}_2) (\mathbf{n}_{12} \cdot \mathbf{p}_1)}{r_{12}} \\ & - \frac{19m_2 + 2\mu - 8m}{4m^2} \frac{(\mathbf{n}_{12} \cdot \mathbf{p}_2 \times \mathbf{S}_1) (\mathbf{n}_{12} \cdot \mathbf{p}_1)}{r_{12}} \\ & \left. \left. - \frac{m_2 + 2\mu - 6m}{2m_1 m} \frac{(\mathbf{p}_1 \cdot \mathbf{p}_2 \times \mathbf{S}_1)}{r_{12}} \right] \right\}. \quad (27) \end{aligned}$$

VI. CONCLUSIONS

We have derived the center-of-mass and rest-frame coordinate representation of the presently known binary Hamiltonians with spinning components. The result is given by

Eq. (17) and Eqs. (25). Additionally, and even more important, all the transformations from the original canonical coordinates to the new ones are presented in explicit form in the Eqs. (19), (20), (26), and (27). This shows full consistency of our general relativistic Hamiltonians having to be of the form (18) with reduced rest-mass only depending on rest-frame coordinates. Thus, our curved spacetimes generated through interacting spinning bodies perfectly respect the asymptotic Lorentz invariance as they should.

Acknowledgments

The authors thank Jan Steinhoff for useful discussions. This work is supported by the Deutsche Forschungsgemeinschaft (DFG) through SFB/TR7 “Gravitational Wave Astronomy” and the DFG project “Krümmungsmaße für Fraktale und Anwendung in der Mustererkennung”.

-
- [1] P. Jaranowski and G. Schäfer, Phys. Rev. D **57**, 7274 (1998).
 - [2] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **62**, 021501 (2000).
 - [3] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **63**, 029903 (2001).
 - [4] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Lett. B **513**, 147 (2001).
 - [5] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **77**, 064032 (2008).
 - [6] J. Steinhoff, G. Schäfer, and S. Hergt, Phys. Rev. D **77**, 104018 (2008).
 - [7] S. Hergt and G. Schäfer, Phys. Rev. D **78**, 124004 (2008).
 - [8] S. Hergt and G. Schäfer, Phys. Rev. D **77**, 104001 (2008).
 - [9] J. Steinhoff, G. Hergt, and S. Schäfer, Phys. Rev. D **78**, 101503(R) (2008).
 - [10] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **62**, 044024 (2000).
 - [11] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **62**, 084011 (2000).
 - [12] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **78**, 024009 (2008).
 - [13] T. J. Rothe, *Grenzfälle des post-newtonschen Dreikörperproblems*, Diploma Thesis in Physics, Friedrich-Schiller-Universität Jena, Jena 2009, unpublished.
 - [14] D. Alba, H. W. Crater, and L. Lusanna, J. Phys. A **40**, 9585 (2007).
 - [15] L. H. Thomas, Phys. Rev. **85**, 868 (1952).

- [16] B. Bakamjian and L. H. Thomas, Phys. Rev. **92**, 1300 (1953).
- [17] L. L. Foldy, Phys. Rev. **122**, 275 (1961).
- [18] J. Stachel and P. Havas, Phys. Rev. D **13**, 1598 (1976).
- [19] M. Pauri and G. M. Prosperi, J. Math. Phys. **17**, 1468 (1976).
- [20] T. D. Newton and E. P. Wigner, Rev. Mod. Phys. **21**, 400 (1949).
- [21] V. A. Brumberg, *Essential Relativistic Celestial Mechanics*, Adam Hilger, Bristol 1991.
- [22] V. M. Vinogradov, Teor. Mat. Fiz. **10**, 338 (1972).
- [23] A. Galindo, Nuovo Cimento **37**, 413 (1965).
- [24] R. N. Hill, J. Math. Phys. **8**, 1756 (1967).
- [25] M. Lorente and P. Roman, J. Math. Phys. **15**, 70 (1974).
- [26] J. Steinhoff and H. Wang, Phys. Rev. D, **81**, 024022 (2010).
- [27] J. M. Martín-García, Computer Phys. Comm. **179**, 597 (2008).
- [28] G. Schäfer, *Post-Newtonian methods: Analytic results on the binary problem* (Proceedings of the CNRS School on Mass, Orléans/France, 2008), arXiv:0910.2857.

Appendix A: Generators given in center-of-mass coordinates

For the convenience of the reader, we also give the resulting generating functions in center-of-mass coordinates. The canonical transformation generated by $-g_{\text{point}}$ or $-g_{\text{spin}}^{\text{total}}$ respectively maps the center-of-mass coordinates onto their Newtonian counterparts:

$$-g_{\text{point}} = h_{1\text{pN}} + h_{2\text{pN}} + h_{3\text{pN}}^{0\text{G}} + h_{3\text{pN}}^{1\text{G}} + h_{3\text{pN}}^{2\text{G}} + h_{3\text{pN}}^{3\text{G}} + O(c^{-8}), \quad (\text{A1})$$

$$-g_{\text{spin}}^{\text{total}} = h_{1\text{pN}} + h_{2\text{pN}} + \frac{1}{2} [h_{\text{spin}} + (1 \leftrightarrow 2)] + O(c^{-6}), \quad (\text{A2})$$

$$\begin{aligned} h_{1\text{pN}} = c^{-2} & \left\{ -\frac{1}{2m^2} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p}) r + \frac{m - 2m_2}{2\mu m^2} (\mathbf{n} \cdot \mathbf{P}) p^2 r \right\} \\ & + c^{-2} G\mu \left(\frac{m_2}{m} - \frac{1}{2} \right) (\mathbf{n} \cdot \mathbf{P}), \end{aligned} \quad (\text{A3a})$$

$$\begin{aligned}
h_{2\text{pN}} = & c^{-4} \left\{ \frac{1}{4m^4} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p}) P^2 r + \frac{m_2 - m_1}{4\mu m^4} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p})^2 r + \frac{m_2 - m_1}{8\mu m^4} (\mathbf{n} \cdot \mathbf{P}) p^2 P^2 r \right. \\
& + \frac{m - 2\mu}{4\mu^2 m^3} (\mathbf{n} \cdot \mathbf{P}) p^2 (\mathbf{P} \cdot \mathbf{p}) r - \frac{m - 2m_2}{8\mu^3 m^2} (\mathbf{n} \cdot \mathbf{P}) (p^2)^2 r \left. \right\} \\
& + c^{-4} G \left\{ \frac{\mu(m - 2m_2)}{8m^3} (\mathbf{n} \cdot \mathbf{P})^3 - \frac{m - 4\mu}{8m^2} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{n} \cdot \mathbf{P})^2 + \frac{m_2 - m_1}{4m^2} (\mathbf{n} \cdot \mathbf{p})^2 (\mathbf{n} \cdot \mathbf{P}) \right. \\
& + \frac{\mu(m - 2m_2)}{8m^3} (\mathbf{n} \cdot \mathbf{P}) P^2 - \frac{5}{8m} (\mathbf{n} \cdot \mathbf{p}) P^2 + \frac{1}{m} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p}) \\
& \left. + \frac{(2\mu - 5m)(m - 2m_2)}{4\mu m^2} (\mathbf{n} \cdot \mathbf{P}) p^2 \right\} - c^{-4} G^2 \frac{\mu(2\mu - m)(m - 2m_2)}{4m} \frac{(\mathbf{n} \cdot \mathbf{P})}{r}, \quad (\text{A3b})
\end{aligned}$$

$$\begin{aligned}
h_{3\text{pN}}^{0\text{G}} = & c^{-6} \left\{ -\frac{1}{6m^6} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p}) (P^2)^2 r + \frac{m - 2m_2}{4\mu m^6} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p})^2 P^2 r \right. \\
& + \frac{5\mu - 2m}{12\mu^2 m^5} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p})^3 r + \frac{7(m - 2m_2)}{96\mu m^6} (\mathbf{n} \cdot \mathbf{P}) p^2 (P^2)^2 r \\
& + \frac{8\mu - 5m}{24\mu^2 m^5} (\mathbf{n} \cdot \mathbf{P}) p^2 (\mathbf{P} \cdot \mathbf{p}) P^2 r - \frac{(4\mu - 7m)(m - 2m_2)}{24\mu^3 m^5} (\mathbf{n} \cdot \mathbf{P}) p^2 (\mathbf{P} \cdot \mathbf{p})^2 r \\
& - \frac{(2m_2 - m)(4\mu + 5m)}{96\mu^3 m^5} (\mathbf{n} \cdot \mathbf{P}) (p^2)^2 P^2 r + \frac{10\mu - 3m}{16\mu^4 m^3} (\mathbf{n} \cdot \mathbf{P}) (p^2)^2 (\mathbf{P} \cdot \mathbf{p}) r \\
& \left. + \frac{m_1^4 - m_2^4}{16\mu^5 m^5} (\mathbf{n} \cdot \mathbf{P}) (p^2)^3 r \right\}, \quad (\text{A3c})
\end{aligned}$$

$$\begin{aligned}
h_{3\text{pN}}^{1\text{G}} = & c^{-6} G \left\{ \frac{m_2(m_2 - m_1)m_1}{32m^6} (\mathbf{n} \cdot \mathbf{P})^5 + \frac{m - 4\mu}{32m^4} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{n} \cdot \mathbf{P})^4 \right. \\
& + \frac{3(m - 2m_2)}{16m^4} (\mathbf{n} \cdot \mathbf{p})^2 (\mathbf{n} \cdot \mathbf{P})^3 - \frac{3(m - 4\mu)}{16\mu m^3} (\mathbf{n} \cdot \mathbf{p})^3 (\mathbf{n} \cdot \mathbf{P})^2 \\
& - \frac{3(m_1 - m_2)}{16\mu m^3} (\mathbf{n} \cdot \mathbf{p})^4 (\mathbf{n} \cdot \mathbf{P}) + \frac{m_2(m_2 - m_1)m_1}{12m^6} (\mathbf{n} \cdot \mathbf{P})^3 P^2 \\
& + \frac{3m - 8\mu}{32m^4} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{n} \cdot \mathbf{P})^2 P^2 - \frac{(2m_2 - m)(\mu + 2m)}{16\mu m^4} (\mathbf{n} \cdot \mathbf{p})^2 (\mathbf{n} \cdot \mathbf{P}) P^2 \\
& - \frac{3}{16\mu m^2} (\mathbf{n} \cdot \mathbf{p})^3 P^2 + \frac{7m_2 m_1 (m_2 - m_1)}{96m^6} (\mathbf{n} \cdot \mathbf{P}) (P^2)^2 \\
& + \frac{1}{4m^3} (\mathbf{n} \cdot \mathbf{p}) (P^2)^2 + \frac{5(m - 4\mu)}{48m^4} (\mathbf{n} \cdot \mathbf{P})^3 (\mathbf{P} \cdot \mathbf{p}) \\
& + \frac{(2\mu - 3m)(m - 2m_2)}{8\mu m^4} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{n} \cdot \mathbf{P})^2 (\mathbf{P} \cdot \mathbf{p}) \\
& + \frac{1}{4\mu m^2} (\mathbf{n} \cdot \mathbf{p})^2 (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p}) + \frac{4\mu - 13m}{24m^4} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p}) P^2 \\
& + \frac{m_2 - m_1}{8\mu m^3} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{P} \cdot \mathbf{p}) P^2 - \frac{(8\mu - 29m)(m - 2m_2)}{24\mu m^4} (\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p})^2 \\
& - \frac{1}{8\mu m^2} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{P} \cdot \mathbf{p})^2 - \frac{(4\mu - 13m)(m - 2m_2)}{96\mu m^4} (\mathbf{n} \cdot \mathbf{P})^3 p^2 \\
& - \frac{1}{2m^3} (\mathbf{n} \cdot \mathbf{p}) (\mathbf{n} \cdot \mathbf{P})^2 p^2 + \frac{m - 2m_2}{4\mu m^3} (\mathbf{n} \cdot \mathbf{p})^2 (\mathbf{n} \cdot \mathbf{P}) p^2 \\
& - \frac{(32\mu - 17m)(m - 2m_2)}{96\mu m^4} (\mathbf{n} \cdot \mathbf{P}) p^2 P^2 \\
& + \frac{1}{4\mu m^2} (\mathbf{n} \cdot \mathbf{p}) p^2 P^2 - \frac{8\mu^2 - 26m\mu + 9m^2}{8\mu^2 m^3} (\mathbf{n} \cdot \mathbf{P}) p^2 (\mathbf{P} \cdot \mathbf{p}) \\
& \left. + \frac{9}{16} \left(\frac{1}{m_2^3} - \frac{1}{m_1^3} \right) (\mathbf{n} \cdot \mathbf{P}) (p^2)^2 \right\}, \tag{A3d}
\end{aligned}$$

$$\begin{aligned}
h_{3\text{pN}}^{2\text{G}} = & c^{-6} G^2 \left\{ \frac{\mu(6\mu - 5m)(m - 2m_2)}{24m^3} \frac{(\mathbf{n} \cdot \mathbf{P})^3}{r} + \left(\frac{\mu^2}{m^2} - \frac{3\mu}{4m} + \frac{5}{48} \right) \frac{(\mathbf{n} \cdot \mathbf{p}) (\mathbf{n} \cdot \mathbf{P})^2}{r} \right. \\
& + \frac{(m - 2m_2)(m - 2\mu)}{4m^2} \frac{(\mathbf{n} \cdot \mathbf{p})^2 (\mathbf{n} \cdot \mathbf{P})}{r} + \frac{\mu(7\mu - 4m)(m - 2m_2)}{24m^3} \frac{(\mathbf{n} \cdot \mathbf{P}) P^2}{r} \\
& + \left(\frac{65}{48} - \frac{7\mu}{8m} \right) \frac{(\mathbf{n} \cdot \mathbf{p}) P^2}{r} + \left(\frac{\mu^2}{2m^2} - \frac{9\mu}{4m} - \frac{101}{48} \right) \frac{(\mathbf{n} \cdot \mathbf{P}) (\mathbf{P} \cdot \mathbf{p})}{r} \\
& \left. + \frac{m - 2m_2}{4m} \frac{(\mathbf{n} \cdot \mathbf{p}) (\mathbf{P} \cdot \mathbf{p})}{r} - \frac{(2m_2 - m)(8\mu^2 + 5m\mu + 30m^2)}{16\mu m^2} \frac{(\mathbf{n} \cdot \mathbf{P}) p^2}{r} \right\}, \tag{A3e}
\end{aligned}$$

$$h_{3\text{pN}}^{3\text{G}} = c^{-6} G^3 \frac{\mu(2m_2 - m)(4\mu^2 - m\mu + m^2)}{8m} \frac{(\mathbf{n} \cdot \mathbf{P})}{r^2}, \tag{A3f}$$

$$\begin{aligned}
h_{\text{spin}} = & c^{-2} \frac{-1}{m_1 m} (\mathbf{P} \cdot \mathbf{p} \times \mathbf{S}_1) + c^{-4} \left\{ \frac{1}{4m_1 m^3} (\mathbf{P} \cdot \mathbf{p} \times \mathbf{S}_1) P^2 + \frac{1}{4m_1^2 m^2} (\mathbf{P} \cdot \mathbf{p} \times \mathbf{S}_1) (\mathbf{P} \cdot \mathbf{p}) \right. \\
& + \left. \frac{m_2 + 2m_1}{4m_2 m_1^3 m} (\mathbf{P} \cdot \mathbf{p} \times \mathbf{S}_1) p^2 \right\} + c^{-4} G \left\{ - \frac{3(m_2 - 2\mu)}{2m_1 m} \frac{(\mathbf{n} \cdot \mathbf{P}) (\mathbf{n} \cdot \mathbf{S}_1)^2}{r^2} \right. \\
& + \frac{3(m - 2m_2)}{2m^2} \frac{(\mathbf{n} \cdot \mathbf{P}) (\mathbf{n} \cdot \mathbf{S}_2) (\mathbf{n} \cdot \mathbf{S}_1)}{r^2} + \frac{4m_2}{m_1 m} \frac{(\mathbf{n} \cdot \mathbf{S}_1) (\mathbf{P} \cdot \mathbf{S}_1)}{r^2} - \frac{1}{m} \frac{(\mathbf{n} \cdot \mathbf{S}_2) (\mathbf{P} \cdot \mathbf{S}_1)}{r^2} \\
& - \frac{3m_2 + 2\mu}{2m_1 m} \frac{(\mathbf{n} \cdot \mathbf{P}) \mathcal{S}_1^2}{r^2} + \frac{m_2 - m_1}{2m^2} \frac{(\mathbf{n} \cdot \mathbf{P}) (\mathbf{S}_1 \cdot \mathbf{S}_2)}{r^2} - \frac{3m_2 + 2\mu}{4m^2} \frac{(\mathbf{n} \cdot \mathbf{P} \times \mathbf{S}_1) (\mathbf{n} \cdot \mathbf{P})}{r} \\
& + \frac{1}{m} \frac{(\mathbf{n} \cdot \mathbf{P} \times \mathbf{S}_1) (\mathbf{n} \cdot \mathbf{p})}{r} - \frac{9m_2 + 2\mu - 4m}{2m_1 m} \frac{(\mathbf{n} \cdot \mathbf{p} \times \mathbf{S}_1) (\mathbf{n} \cdot \mathbf{P})}{r} \\
& \left. - \frac{m_2 + 2\mu - 6m}{2m_1 m} \frac{(\mathbf{P} \cdot \mathbf{p} \times \mathbf{S}_1)}{r} \right\}. \tag{A4}
\end{aligned}$$